

# Administration of CD4<sup>+</sup>CD25<sup>high</sup>CD127<sup>-</sup> Regulatory T Cells Preserves $\beta$ -Cell Function in Type 1 Diabetes in Children

NATALIA MAREK-TRZONKOWSKA, VMD, PHD<sup>1</sup>  
 MAŁGORZATA MYŚLIWIEC, MD, PHD<sup>2</sup>  
 ANITA DOBYSZUK, MSc<sup>1</sup>  
 MARCELINA GRABOWSKA, MSc<sup>1</sup>  
 ILONA TECHMAŃSKA, MD<sup>2</sup>  
 JOLANTA JUŚCIŃSKA, MD, PHD<sup>3</sup>

MAGDALENA A. WUJTEWICZ, MD, PHD<sup>4</sup>  
 PIOTR WITKOWSKI, MD, PHD<sup>5</sup>  
 WOJCIECH MŁYNARSKI, MD, PHD<sup>6</sup>  
 ANNA BALCERSKA, MD, PHD<sup>2</sup>  
 JOLANTA MYŚLIWSKA, MD, PHD<sup>7</sup>  
 PIOTR TRZONKOWSKI, MD, PHD<sup>1,8</sup>

**OBJECTIVE**—Type 1 diabetes is a condition in which pancreatic islets are destroyed by self-reactive T cells. The process is facilitated by deficits in the number and suppressive activity of regulatory T cells (Tregs). Here, we show for the first time that the infusion of autologous Tregs prolongs remission in recently diagnosed type 1 diabetes in children.

**RESEARCH DESIGN AND METHODS**—We have administered Tregs in 10 type 1 diabetic children (aged 8–16 years) within 2 months since diagnosis. In total, 4 patients received  $10 \times 10^6$  Tregs/kg body wt, and the remaining 6 patients received  $20 \times 10^6$  Tregs/kg body wt. The preparation consisted of sorted autologous CD3<sup>+</sup>CD4<sup>+</sup>CD25<sup>high</sup>CD127<sup>-</sup> Tregs expanded under good manufacturing practice conditions.

**RESULTS**—No toxicity of the therapy was noted. A significant increase in the percentage of Tregs in the peripheral blood has been observed since the day of infusion. These patients were followed along with matched type 1 diabetic patients not treated with Tregs. Half a year after type 1 diabetes onset (4–5 months after Tregs infusion), 8 patients treated with Tregs still required  $<0.5$  UI/kg body wt of insulin daily, with 2 patients out of insulin completely, whereas the remission was over in the nontreated group. In addition, plasma C-peptide levels were significantly higher in the treated group as compared with those not treated.

**CONCLUSIONS**—This study shows that the administration of Tregs is safe and tolerable in children with recent-onset type 1 diabetes.

*Diabetes Care* 35:1817–1820, 2012

Type 1 diabetes is a rising problem around the world. For example, it is estimated that the morbidity in Poland doubles every 10 years (1). The condition develops as a result of autoimmune attack of self-reactive T cells that infiltrate pancreatic islets and destroy insulin-producing

$\beta$ -cells (2,3). The transfer of self-reactive T cells from diabetic animals induces insulinitis and diabetes in previously healthy animals (4). This autoimmune process can be prevented by highly suppressive regulatory T cells (Tregs) (5,6). It has been found in several animal models that CD4<sup>+</sup>CD25<sup>+</sup>FoxP3<sup>+</sup>

Tregs can stop the destruction of pancreatic islets and protect from autoimmune type 1 diabetes (7,8). In humans, mutation in the *foxp3* gene results in immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome, which is associated with lack of functional Tregs, and subsequent autoimmune diabetes (9).

Tregs can regulate ongoing immune reactions and, therefore, can be used to suppress them. In 2009, we successfully administered expanded Tregs in patients with chronic graft-versus-host disease (10). The efficacy of such an approach recently has been confirmed in the prophylaxis of graft-versus-host disease (11,12). Here, we show the effects of adoptive transfer of Tregs in children with recently diagnosed type 1 diabetes.

## RESEARCH DESIGN AND METHODS

In the current study, 10 children (aged 8–16 years, 6 girls and 4 boys) with type 1 diabetes, diagnosed in compliance with World Health Organization criteria (13), were treated with Tregs (Supplementary Fig. 1). The inclusion criteria were age 5–18 years; up to 2 months since type 1 diabetes diagnosis; the presence of anti-GAD65 antibody, islet cell antibody, and insulin autoantibody; fasting C-peptide levels  $>0.4$  ng/mL in two consecutive measurements (a challenge test was not permitted by the ethics committee); BMI range of 25th–75th percentiles for a particular age; and adequate venous access. The following exclusion criteria were applied: any cytopenia or low hemoglobin levels; carriage of HLA-DQB1\*0602 allele; positive test for hepatitis B virus, hepatitis C virus, HIV, *Treponema pallidum*, or other active infections; history of neoplasm; excessive anxiety related to the procedure; and chronic disease requiring pharmacological treatment. Up to 250 mL peripheral blood was drawn with anesthesiological assistance. For children with body weight  $<50$  kg, the blood volume drawn was equal to 0.5% of body weight (see study protocol and patient consent form in the Supplementary Data online).

Tregs were prepared as previously described (10,14). In brief, the blood cells

From the <sup>1</sup>Department of Clinical Immunology and Transplantology, Medical University of Gdańsk, Gdańsk, Poland; the <sup>2</sup>Department of Pediatrics, Hematology, Oncology, and Endocrinology, Medical University of Gdańsk, Gdańsk, Poland; the <sup>3</sup>Blood Bank of the Pomerania Region, Gdańsk Unit, Gdańsk, Poland; the <sup>4</sup>Department of Anesthesiology and Critical Care, Medical University of Gdańsk, Gdańsk, Poland; the <sup>5</sup>Department of Surgery, Section of Transplantation, University of Chicago, Chicago, Illinois; the <sup>6</sup>Department of Pediatrics, Oncology, Hematology, and Diabetology, Medical University of Lodz, Lodz, Poland; the <sup>7</sup>Department of Immunology, Medical University of Gdańsk, Gdańsk, Poland; and the <sup>8</sup>Tricity Academic Experimental Facility, Medical University of Gdańsk, Gdańsk, Poland.

Corresponding author: Piotr Trzonkowski, ptrzon@gumed.edu.pl.

Received 6 January 2012 and accepted 3 April 2012.

DOI: 10.2337/dc12-0038. Clinical trial reg. no. NKEBN/8/2010 (phase 1).

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc12-0038/-DC1>.

A slide set summarizing this article is available online.


N.M.-T. and M.M. contributed equally to this study.

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

were sorted to CD3<sup>+</sup>CD4<sup>+</sup>CD25<sup>high</sup>CD127<sup>-</sup>lin<sup>-</sup>doublet<sup>-</sup> Tregs using combined immunomagnetic/fluorescence-activated technique with the purity of ~100% (median [minimum–maximum] 98% [97–99]). As compared with our previous protocol, important improvements in cell processing were the application of the good manufacturing practice (GMP)-adapted fluorescence-activated cell sorter (FACS) equipped with single-use sample lines (Influx; BD Biosciences, San Jose, CA), which eliminated the risk of cross-contamination, and the application of GMP-grade CellGro medium (CellGenix, Breisgau, Germany) supplemented with 10% autologous serum, interleukin-2 (1,000 units/mL; Proleukin; Chiron, San Diego, CA), and clinical-grade anti-CD3/anti-CD28 beads (Invitrogen, Carlsbad, CA) in 1:1 ratio with cells. Tregs were cultured until the required number was achieved, but for no longer than 2 weeks (median [minimum–maximum]: 10 days [7–14]). Taking all these measures into consideration, we have significantly improved the stability and quality of the Tregs in the final product. The release criteria for Tregs preparation were FoxP3 expression >90% (93% [90–97]), passed interferon (IFN)- $\gamma$  suppression assay, and negative microbiological tests (no genetic material of hepatitis B virus, hepatitis C virus, or HIV and no microbial contamination in the culture supernatants). Tregs for infusion were washed out completely, suspended in 250 mL 0.9% NaCl (Polfa, Starogard Gd., Poland), and transferred in slow infusion to the patient under anesthesiological

assistance within 1 h of product release. The administered dose was either 20  $\times$  10<sup>6</sup>/kg body wt (n = 6) or when protocol required cessation of Tregs expansion at 2 weeks, 10  $\times$  10<sup>6</sup>/kg body wt (n = 4). As control, the other 10 type 1 diabetic patients and their respective matching characteristics were included in the follow-up but did not receive Tregs. The trial was neither blind nor randomized, and the comparison group did not undergo any interventions, such as initial blood drawing for cell processing or mock transfusions (comparison of clinical characteristics of the groups provided in Table 1). End points of the study were fasting C-peptide, HbA<sub>1c</sub> levels, and insulin requirement, with special attention to daily insulin dose (DDI; 0.5 UI/kg body wt) as a borderline for remission in our protocol. The study was conducted according to the protocol approved by the Ethics Committee of the Medical University of Gdańsk (NKEBN/8/2010). Written informed consent was obtained before starting the procedures.

**RESULTS**—No serious infections, episodes of acute hyper- or hypoglycemia, or other adverse effects were observed after the administration of Tregs during the entire follow-up. In one of the patients, the infusion was coincident with influenza, which was diagnosed a day after the procedure (confirmed from nasopharyngeal specimen [bioNexia Influenza; bioMérieux, Marcy l’Etoile, France]). The administration of Tregs was followed by a significant decrease in the requirement of exogenous insulin and a decrease in HbA<sub>1c</sub> levels in all



**Figure 1**—C-peptide, DDI, HbA<sub>1c</sub>, and fasting glucose in type 1 diabetic children treated with Tregs infusions (n = 10) during follow-up. Values for matched comparison patients not treated with Tregs (n = 10) are shown as gray columns. Values are given as median (minimum–maximum). \*Significant differences.


the patients after 2 weeks (Fig. 1 and Supplementary Figs. 2 and 3). In addition, a significant increase in the percentage of Tregs in the peripheral blood was observed since the day of treatment (Wilcoxon test, P = 0.04) (Fig. 2 and Supplementary Figs. 2 and 3).

Significant differences between the study group and the comparison group were noted for the first time half a year after the diagnosis of type 1 diabetes (4–5 months after Tregs administration). Of the 10 patients treated with Tregs, 8 were still in clinical remission, defined as DDI <0.5 UI/kg body wt (DDI median [minimum–maximum] 0.22 UI/kg body wt [0–0.60]),

**Table 1**—Clinical characteristics of the patients

	Treated (n = 10)	Not treated (n = 10)
Age (years)	12.2 (8–16)	11.8 (7–16)
BMI	16.9 (14.2–21.5)	16.9 (14.2–20.7)
Fasting glycemia at diagnosis (mg%)	354 (151–588)	354 (151–598)
Polydipsia at diagnosis	5	8
Polyuria at diagnosis	5	8
Loss of weight at diagnosis	4	3
pH at diagnosis (capillary blood)	7.39 (7.36–7.46)	7.39 (7.34–7.53)
pO <sub>2</sub> at diagnosis (capillary blood, mmHg)	69.3 (63.4–88.0)	69.0 (56.9–86.2)
pCO <sub>2</sub> at diagnosis (capillary blood, mmHg)	39.1 (28.0–41.8)	38.0 (24.0–41.0)
HCO <sub>3</sub> at diagnosis (capillary blood, mmHg)	23.85 (18.8–25.0)	23.3 (21.3–25.2)
Acid/base balance at diagnosis (BE, mEq/L)	–0.55 (–7.8 to 1.0)	–0.6 (–7.0 to 0.9)
SatO <sub>2</sub> at diagnosis (capillary blood, %)	94.1 (90.2–97.3)	95.5 (92.4–98.0)
Anti-GAD65 antibody	10	10
Islet cell antibody	5	5
Insulin autoantibody	9	8

Data are median (minimum–maximum) or number of patients. BE, base excess; pCO<sub>2</sub>, pressure of carbon dioxide; pO<sub>2</sub>, pressure of oxygen; SatO<sub>2</sub>, oxygen saturation.



**Figure 2**—Levels of  $CD3^+CD4^+CD25^{high}CD127^{low}FoxP3^+$  Tregs in type 1 diabetic children treated with Tregs infusions ( $n = 10$ ) during follow-up. Value for the  $-10$ -days point refers to the day of blood drawing for Tregs sorting. Gray column depicts values measured for the matched comparison patients not treated with Tregs (only six patients were available). Data are shown as medians (minimum–maximum). \*Significant differences.

with 2 patients out of insulin completely. At the same time, remission was completed in 6 of 10 patients in the comparison group not treated with Tregs (DDI 0.52 U/kg body wt [0.18–0.69]; Mann-Whitney  $U$  test,  $P = 0.04$ ). In addition, plasma C-peptide levels were significantly higher in patients from the Tregs group as compared with the untreated group (0.75 [0.38–2.11] vs. 0.48 ng/mL [0.15–0.56]; Mann-Whitney  $U$  test,  $P = 0.01$ ) (Fig. 1). No differences were seen between the two doses of Tregs used and, therefore, patients are presented as a single cohort.

**CONCLUSIONS**—As previously described, Tregs from diabetic patients can be effectively expanded (15). However, there were doubts whether autologous Tregs could be effective in the treatment of type 1 diabetes. It is notable that a fraction of  $IFN-\gamma^+FoxP3^+$  Tregs with reduced suppressive efficiency have been described in diabetic patients (16). We could also detect such cells, albeit in a very low proportion in our type 1 diabetic children ( $\sim 1\%$  of Tregs; data not shown), and they have never affected the results of the suppression assay in vitro. Hence, it might be possible that the majority of defects in type 1 diabetes are related to self-reactive effector T cells, which might be difficult to control by Tregs (17,18). Such activated effector T cells specific for self-islets have been documented recently in humans (19). Fortunately, in NOD mice, the transfer of Tregs could suppress the activity of self-reactive effector T cells and stop diabetes (20). Our study

confirms a similar effect of such a Tregs transfer in the human setting.

The dose of Tregs administered in our study has a biological meaning because it was able to significantly increase the level of  $FoxP3^+$  Tregs in peripheral blood. There were several issues that made our protocol efficacious. In all published trials, immunomagnetic isolation of Tregs was applied, resulting in a maximum of  $\sim 60\%$   $FoxP3^+$  T cells in the product, but that figure could be  $<20\%$  (11,12). So, at least 40% of cells in the product administered to the patients were not Tregs from the beginning. This made it difficult to assess the exact level of administered cells with immunosuppressive activity. With FACS sorting, we reliably achieved  $\sim 100\%$   $CD3^+CD4^+CD25^{high}CD127^{low}FoxP3^+$  T cells, and in children  $>90\%$  of those cells were still  $FoxP3^+$  Tregs after expansion. Although FACS sorting is considered a non-GMP procedure, its safety with new sorters is comparable with that of clinical-grade immunomagnetic sorting. In fact, novel single-use fluidic lines preventing cross-contamination and performing the procedure in clean room facilities, together with the resulting purity, make FACS sorting superior to immunomagnetic sorting. Another problem somehow related to the dose assessment is a quality check of the ready-to-administer expanded Tregs product. Proposed in the literature, suppression of the proliferation assay could not be included in release criteria because the result was available 5 days after commencing the assay. In this time period, the suppression of expanded Tregs could change dramatically. Our  $IFN-\gamma$  suppression assay could be included in release criteria, since the results were available within hours, and suppressive activity of the product was confirmed prior to the transfer of Tregs to patients.

There are several limitations of the study involving number of patients and timing. In addition, it is probable that the fast initial decrease in the level of  $HbA_{1c}$  was very much related to the improvement of the clinical status of the patients and medical interventions other than Tregs infusion. Nevertheless, even at this early 4-month point of our study, when we compare our patients with carefully matched untreated subjects, we already see the difference in the insulin dose and C-peptide levels. Further follow-up, which is under way, will confirm whether this is a sustained remission or only a delay in the destruction of islets.

In our opinion, maneuvers increasing the number of Tregs are safe and tolerable

and may potentially control type 1 diabetes. As this study shows, in children with recently diagnosed type 1 diabetes, some improvement in the functioning of  $\beta$ -cells after the infusion of expanded autologous Tregs is a proof of this concept. These results should encourage further investigations involving our novel approach.

**Acknowledgments**—The study was supported by the Polish Ministry of Science (grants NN402-3530-38 and NR13-0126-10). N.M.-T. is a Fellow of the Kosciuszko Foundation.

No potential conflicts of interest relevant to this article were reported.

N.M.-T., M.M., and P.T. contributed to the study design, protocol writing, cell preparation, data collection, analysis, interpretation, and writing and reviewing of the manuscript. A.D. contributed to cell separation and data collection. M.G. and I.T. contributed to data collection and interpretation. J.J. and M.A.W. contributed to data collection. P.W. contributed to data analysis and interpretation and reviewed the report. W.M. contributed to data collection, analysis, and interpretation and reviewed the manuscript. A.B. and J.M. contributed to data collection and interpretation and reviewed the manuscript. All authors have approved the manuscript for submission. P.T. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Parts of this study were presented in abstract form at the 72nd Scientific Sessions of the American Diabetes Association, Philadelphia, Pennsylvania, 8–12 June 2012.

The authors thank Lucyna Szumacher-Sharma from the Department of Pediatrics, Hematology, Oncology, and Endocrinology, Medical University of Gdańsk, for her perfect assistance in clinical procedures.

## References

- Jaroszc-Chobot P, Polanska J, Szadkowska A, et al. Rapid increase in the incidence of type 1 diabetes in Polish children from 1989 to 2004, and predictions for 2010 to 2025. *Diabetologia* 2011;54:508–515
- Gepts W, Lecompte PM. The pancreatic islets in diabetes. *Am J Med* 1981;70:105–115
- Eisenbarth GS. Type I diabetes mellitus. A chronic autoimmune disease. *N Engl J Med* 1986;314:1360–1368
- Bendelac A, Carnaud C, Boitard C, Bach JF. Syngeneic transfer of autoimmune diabetes from diabetic NOD mice to healthy neonates. Requirement for both  $L3T4^+$  and  $Lyt-2^+$  T cells. *J Exp Med* 1987;166:823–832
- Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor  $\alpha$ -chains (CD25). Breakdown of a single mechanism of self-tolerance causes

- various autoimmune diseases. *J Immunol* 1995;155:1151–1164
6. Salomon B, Lenschow DJ, Rhee L, et al. B7/CD28 costimulation is essential for the homeostasis of the CD4+CD25+ immunoregulatory T cells that control autoimmune diabetes. *Immunity* 2000;12:431–440
  7. Green EA, Gorelik L, McGregor CM, Tran EH, Flavell RA. CD4+CD25+ T regulatory cells control anti-islet CD8+ T cells through TGF- $\beta$ -TGF- $\beta$  receptor interactions in type 1 diabetes. *Proc Natl Acad Sci U S A* 2003;100:10878–10883
  8. You S, Slehoffer G, Barriot S, Bach JF, Chatenoud L. Unique role of CD4+CD62L+ regulatory T cells in the control of autoimmune diabetes in T cell receptor transgenic mice. *Proc Natl Acad Sci U S A* 2004;101 (Suppl. 2):14580–14585
  9. Bennett CL, Christie J, Ramsdell F, et al. The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet* 2001;27:20–21
  10. Trzonkowski P, Bieniaszewska M, Juścińska J, et al. First-in-man clinical results of the treatment of patients with graft versus host disease with human ex vivo expanded CD4+CD25+CD127- T regulatory cells. *Clin Immunol* 2009;133:22–26
  11. Di Ianni M, Falzetti F, Carotti A, et al. Tregs prevent GVHD and promote immune reconstitution in HLA-haploidentical transplantation. *Blood* 2011;117:3921–3928
  12. Brunstein CG, Miller JS, Cao Q, et al. Infusion of ex vivo expanded T regulatory cells in adults transplanted with umbilical cord blood: safety profile and detection kinetics. *Blood* 2011;117:1061–1070
  13. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539–553
  14. Marek N, Bieniaszewska M, Krzystyniak A, et al. The time is crucial for ex vivo expansion of T regulatory cells for therapy. *Cell Transplant* 2011;21:1747–1758
  15. Putnam AL, Brusko TM, Lee MR, et al. Expansion of human regulatory T-cells from patients with type 1 diabetes. *Diabetes* 2009;58:652–662
  16. McClymont SA, Putnam AL, Lee MR, et al. Plasticity of human regulatory T cells in healthy subjects and patients with type 1 diabetes. *J Immunol* 2011;186:3918–3926
  17. D'Alise AM, Auyeung V, Feuerer M, et al. The defect in T-cell regulation in NOD mice is an effect on the T-cell effectors. *Proc Natl Acad Sci U S A* 2008;105:19857–19862
  18. Schneider A, Rieck M, Sanda S, Pihoker C, Greenbaum C, Buckner JH. The effector T cells of diabetic subjects are resistant to regulation via CD4+ FOXP3+ regulatory T cells. *J Immunol* 2008;181:7350–7355
  19. Velthuis JH, Unger WW, Abreu JR, et al. Simultaneous detection of circulating autoreactive CD8+ T-cells specific for different islet cell-associated epitopes using combinatorial MHC multimers. *Diabetes* 2010;59:1721–1730
  20. Bluestone JA, Tang Q. Therapeutic vaccination using CD4+CD25+ antigen-specific regulatory T cells. *Proc Natl Acad Sci U S A* 2004;101 (Suppl. 2):14622–14626